

*Stenichnus* 属の種は、日本からは北海道と九州、トカラ列島及び八重山諸島から 5 種類が確認されているのみで、本州からは未記録であったが、本種はこの分布の間を埋める種となる。

### References

- JALOSZYŃSKI, P., 2004. Revision of *Stenichnus* THOMSON (Coleoptera, Scydmaenidae) of Japan and Taiwan. *Bull. Nat. Sci. Mus. Tokyo*, **30**(4): 155–171.
- 2006. A new species of *Stenichnus* THOMSON (Coleoptera, Scydmaenidae) from Nakanoshima Is., Tokara Archipelago, Japan. *Bull. Nat. Sci. Mus. Tokyo*, **32**(2): 53–56.

---

*Elytra, Tokyo*, **37**(1): 69–70, May 29, 2009

## Rediscovery of *Stenhomalus japonicus* (Coleoptera, Cerambycidae) from Island of Sado of Type Locality

Tatsuya NIISATO<sup>1)</sup> and Tomio KINOSHITA<sup>2)</sup>


<sup>1)</sup> Bioindicator Co., Ltd., Yarai-chô 126, Shinjuku, Tokyo, 185–0206 Japan

<sup>2)</sup> Kamisôyagi 8–20–6, Yamato City, Kanagawa, 242–0029 Japan

*Stenhomalus japonicus* (PIC, 1904) is one of the most problematical species among the Japanese members of the Obriini. This species had been believed as a member of the genus *Obrium* followed the original combination, and was recently identified its true systematic position as the result of reexamination of holotype (NIISATO, 2006). Besides, the long used name *S. lighti* GRESSITT, 1935 was suggested as a junior synonym of this species (NIISATO, *op. cit.*). Though originally described from “Sado, Japan”, *S. japonicus* has never been rediscovered from the type locality after the original description. We are going to give an additional record of this species from Sado Island based on the specimens recently collected by the junior author.

*Specimens examined.* 2 ♂♂, 2 ♀♀, Mt. Donden-yama, Sado Is., off NW. Honshu, Ryôtsu City, Niigata Pref., Japan, emerged out in IV–1999, from the freshly dead trunk of *Cornus controversa* HEMSL., T. KINOSHITA leg.

*Notes.* Male genital organs of *S. japonicus* from Sado Island are as shown in Figs. 1–3. MIROSHIKOV (1989) described and illustrated the male genitalia of *S. japonicus* based on the specimen from Far East Russia under the name of *S. lighti*. According to the present examination, the median lobe and vestigial tegmen at least in proportion are quite identical with those of two localities.



Figs. 1–3. Male genital organs of *Stenhomalus japonicus* (PIC) from Sado Is., off NW. Honshu, Japan. — 1. Median lobe with vestigial tegmen, lateral view; 2, ditto, dorsal view; 3, 8th and 9th abdominal segments, ventral view.

### References

- MIROSHIKOV, A. I., 1989. New and little known longicorn beetles (Coleoptera, Cerambycidae) from Far East and the systematic position of the genus *Stenhomalus* WHITE, 1855. *Ent. Obozr.*, **68**: 739–746. (In Russian, with English summary.)
- NIISATO, T., 2006. Taxonomic disorder of *Obrium japonicum* (Coleoptera, Cerambycidae) and its allied species. *Elytra, Tokyo*, **34**: 379–395.
- PIC, M., 1904. Description d'un *Obrium* du Japon et note de chasse. *Mat. Longic.*, **5**(1): 22.